

(Check) 9 - 2016 Jurnal Teknologi(AR2BIO) Nesti Sianipar - Induction, Multiplication and Acclimatization of Red Betel

by Arta Moro Sundjaja

FILE	-		
	_INDUCTION,_MULTIPLICATION_AND_ACCLIMATIZATION_OF_RED_BET EL.PDF (345K)		
TIME SUBMITTED	10-APR-2017 11:49AM	WORD COUNT	4224
SUBMISSION ID	653860491	CHARACTER COUNT	21724

INDUCTION, MULTIPLICATION, AND ACCLIMATIZATION OF RED BETEL PLANT (*PIPER CROCATUM* RUIZ AND PAV.) BY IN VITRO ORGANOGENESIS

Nesti Fronika Sianipara^a, Verlina^b, Rosaria^b

^aFood Technology Department, Faculty of Engineering, Bina Nusantara University, 15325 Tangerang, Indonesia

^bAlumni of Biology Department, Faculty of Science and Technology, Pelita Harapan University, 15811 Tangerang, Indonesia

Article history

Received

25 June 2015

Received in revised form

16 September 2015

Accepted

20 December 2015

*Corresponding author
sianipar@binus.edu/
nestipro@yahoo.com

Graphical abstract



Abstract

Red betel is an herbal plant with anticancer activities. Propagation of red betel by conventional vegetative method has low multiplication factors. The purpose of this research was to increase the propagation rate of red betel by utilizing in vitro culture organogenesis method which was consisted of stages as follows: shoots initiation, shoots multiplication, root induction, and acclimatization. Shoots induction from axillary bud was achieved on MS 3 mgL⁻¹ BAP media. Shoots were multiplied on MS (0, 0.5, 1, 1.5 mgL⁻¹) BAP, (0.01, 0.03, 0.1, 0.3 mgL⁻¹) TDZ, and (0.01, 0.03, 0.5, 0 mgL⁻¹) TDZ combined with 0.5 mgL⁻¹ NAA. Roots were induced on MS (0, 0.5, 1, 1, 5 mgL⁻¹) NAA. The maximum number of shoots (4.4 ± 0.9) was achieved on MS 1 mgL⁻¹ BAP. The maximum number of roots (22.8 ± 9.7) was achieved on MS 0.5 mgL⁻¹ NAA. The highest survival rates of red betel plants during acclimatization and post-acclimatization periods were 88% and 100%, respectively. Red betel had been successfully multiplied by in vitro direct organogenesis.

Keywords: *Piper crocatum* ruiz and pav, in vitro organogenesis, induction, multiplication, acclimatization

© 2016 Penerbit UTM Press. All rights reserved

1.0 INTRODUCTION

Red betel (*Piper crocatum* Ruiz and Pav.) is a medicinal herb which has a high potency for curing various diseases, because it has anticancer [1], antihyperglycemic, antidiabetic [2], antiinflammation [3], and antibacterial activities [4]. Red betel is also effective in curing whiteness [5] and acts as natural antioxidants [6]. Red betel contains a broad array of phytochemical compounds such as alkaloids, flavonoids, tannins, and saponins [3, 7].

Red betel has been cultivated by vegetative methods in its natural habitat, specifically by cutting and grafting methods [8]. However, propagation of red betel plant through vegetative methods has

limitations, such as the short storage period of planting materials, slow rate of plant multiplication, and susceptibility to viral infections [9].

The problems of conventional propagation can be overcome by tissue culture method. Tissue culture is able to produce high-quality planting materials, uniform seeds, and disease-resistant plants rapidly [10]. Direct organogenesis is one of the propagation methods involving the formation of buds or organs directly without passing through the callus. Propagation can be achieved through the formation of shoots from axillary bud explants by breaking the apical dominance of the apical meristem [11], such as in the case of propagation of *Piper nigrum* [12].

Murashige Skoog (MS) is the most commonly used basal media in plant tissue culture because it contains macro- and micronutrient to support optimum plant growth. Composition of plant growth regulators in tissue culture media is an instrumental in morphogenesis. Cytokinins is a class of plant growth regulator which is frequently used in propagating various plants in vitro. Cytokinins regulate the formation of axillary buds and lateral roots [11].

The most common types of cytokinins are 6-Benzyl Amino Purine (BAP) and Thiadiazuron (TDZ). BAP is able to induce the growth of axillary bud from shoot tip culture. Multiplication of shoots, buds, and meristems can be optimized by adjusting the concentration of BAP. Low concentration of TDZ is also able to propagate shoots [11]. In vitro herbal plants were successfully induced and multiplied by using BAP and TDZ. Shoots multiplication of *Talinum triangular* [13], *Gynura pseudochina* (L) DC [14], and *Gloriosa superba* [15] were achieved on media supplemented with BAP. Meanwhile, *Bacopa monnieri* was multiplied on media supplemented with TDZ [16]. Plants of the genus *Piper*, such as black pepper (*Piper nigrum*) and *Piper longum* [17] were also successfully propagated by in vitro culture method [12]. Red betel plantlets were needed to be acclimatized because the success of plant propagation by in vitro method was indicated by the survival rates of plants during acclimatization [10]. Research about in vitro culture of Indonesian red betel plant had never been published before. The specific aim of this research was to obtain the optimal media formulation for in vitro organogenesis of red betel which was consisted of shoots initiation, shoots multiplication, root induction, and acclimatization

2.0 EXPERIMENTAL

2.1 Sterilization of Explants

In vitro red betel plantlets were initiated from a single node axillary bud in Biotechnology Laboratory of Pelita Harapan University. The explants were washed with detergent and rinsed with running water to remove soil particles. The explants were sunk in 0.25 gram of both bactericide and fungicide in 200 ml of water for 30 minutes, 2.25 mgL⁻¹ of antibiotic rifampicin for four hours, 2% of Clorox® bleach for 15 minutes, 1.5% of Clorox® bleach for 15 minutes, 1% of Clorox® bleach for 10 minutes, and 0.1% of HgCl₂ for 10 minutes. The explants were then rinsed twice in sterile water and planted into in vitro plant tissue culture media in order to induce the growth of explants.

2.2 Induction and Initiation of Shoots

Induction was achieved on MS media supplemented with 3 mgL⁻¹ BAP in three months. The parameters recorded in the experiment were the

contamination percentages and the mean numbers of shoots produced per explant. The explants were first initiated on MS media supplemented with 3 mgL⁻¹ BAP for four months. Explants were then initiated for the second time on MS media supplemented with either 3 mgL⁻¹ BAP or 4 mgL⁻¹ BAP. The obtained in vitro shoots were used as explants for the next research stages. The parameters recorded in the experiment were the mean number of shoots and leaves produced per explant in two months.

2.3 Multiplication of Shoots

In vitro shoots were multiplied on MS media supplemented with BAP, TDZ, and TDZ combined with 1-Naphthaleneacetic acid (NAA). Explants were cultured on MS media supplemented with different concentration of BAP (0, 0.5 mgL⁻¹, 1 mgL⁻¹, 1.5 mgL⁻¹), and TDZ (0.01 mgL⁻¹, 0.03 mgL⁻¹, 0.1 mgL⁻¹, 0.3 mgL⁻¹). Explants were also cultured on MS media supplemented with different concentration of TDZ (0.01 mgL⁻¹, 0.03 mgL⁻¹, 0.05 mgL⁻¹, 0 mgL⁻¹) combined with 0.5 mgL⁻¹ NAA. The parameters recorded in the experiment were the mean number of shoots and leaves produced and the mean number of shoot lengths in three months.

2.4 Root Induction

Roots were induced on MS media supplemented with (0 mgL⁻¹, 0.5 mgL⁻¹, 1 mgL⁻¹, 1.5 mgL⁻¹) NAA. All of the experiments were replicated 10 times. The parameters recorded in the experiment were the percentage viable rooted plantlets, time needed for root formation, and the mean number of roots in three months.

2.5 Acclimatization

In vitro propagated plantlets of red betel were acclimatized after three months being cultured in rooting media. The plantlets were washed with running tap water and sunk into the mixture of 0.25 gram bactericide and fungicide in 200 ml of water. The plantlets were then planted into acclimatization media which contained husk and compost (1:1) in plastic bottles. The media were always kept moist in room temperature. The survival rates of plants were recorded for four weeks. Viable plants were then moved into post-acclimatization media composed of a mixture of husk, soil, and compost media (1:1:1) in the green house. The survival rates of plantlets were also recorded for three months.

3.0 RESULTS AND DISCUSSION

3.1 Induction and Initiation of Explants

Multiple shoots were induced from one out of 15 axillary buds (6.67%). Multiple shoots were transferred to the first initiation on MS media supplemented with 3 mgL⁻¹ BAP. The mean number of shoots and leaves were recorded after four months (Table 1). Red betel explants were initiated on the second initiation media, i.e. MS supplemented with 3 mgL⁻¹ BAP and 4 mgL⁻¹ BAP. The highest mean numbers of shoots and leaves were obtained on MS supplemented 3 mgL⁻¹ BAP, with 10.25 shoots and 30.75 leaves, respectively (Table 1, Figure 1a, b)

Table 1 Mean number of Shoots and Leaves at 12th week

First Initiation (month) ^a	Mean Number of Shoots ^c	Mean of Number of Leaves ^c
1	5.33	10.67
2	6.11	13.56
3	7.44	14.89
4	8	16
Second Initiation (mg ⁻¹ L) ^b	Mean Number of shoots ^c	Mean Number of leaves ^c
BAP 3	10.25	30.75
BAP 4	7	22.8

^a The explants were first initiated on MS BAP 3 mg L⁻¹. ^b Explants were then initiated for the second time on MS BAP 3 mg⁻¹ L or BAP 4 mg⁻¹ L. ^c the parameters were observed after 12 weeks.

The Formation of shoots and leaves during the induction and initiation phases was influenced by the concentration of cytokinins in the media. BAP is a cytokinin-type plant growth regulator that stimulates the proliferation of axillary buds if it is supplemented in appropriate concentrations [10, 18]. BAP affects the amount of nitrogen and potassium on MS media which are involved in protein synthesis and cell division [19]. The high concentration of BAP in culture media inhibits the differentiation and multiplication of shoots due to the excessive accumulation of cytokinins in the shoots [20].

3.2 Multiplication of Shoots

Multiplication of red betel in vitro plantlets on MS supplemented with BAP obtained the highest mean number of shoots (4.4 ± 0.9), shoot lengths (2.7 ± 1.4 cm), and leaves (20.2 ± 6.1) on media supplemented with 1 mgL⁻¹ BAP. Tukey test at 5% significance level showed that the mean number of shoots on MS media supplemented with 1 mgL⁻¹ BAP was significantly different from control but not significantly different with MS supplemented with 0.5 mgL⁻¹ BAP and 1.5 mgL⁻¹ BAP (Table 2; Figure 1c).

Table 2 Mean Number of Shoots, Shoot Lengths, and Leaves on MS BAP Media at 12th week

Concentration of BAP (mg ⁻¹ L)	Mean Number of Shoots (mean ± SD) ^a	Mean of Shoot lengths (cm) (mean ± SD) ^a	Mean Number of Leaves (mean ± SD) ^a
0	2.6 ^a ± 0.8	1.5 ^a ± 0.17	13.8 ^{ab} ± 4.9
0.5	3.4 ^{ab} ± 0.9	1.5 ^a ± 0.3	9.2 ^a ± 4.5
1	4.4 ^b ± 0.9	2.7 ^a ± 1.4	20.2 ^b ± 6.1
1.5	3.6 ^{ab} ± 0.5	1.1 ^a ± 0.3	14 ^{ab} ± 1.6

^a Means followed by same letters are not significantly different at p ≤ 0.05 according to Tukey multiple range test

MS basal media supplemented with BAP 1 mgL⁻¹ gave the highest mean number of shoots, shoot lengths, and number of leaves. In this study, it appeared that the control medium (MS) was also able to induce the production of shoots. It was probably due to the endogenous cytokinins already existed in red betel plantlets that triggered cell differentiation and bud formation [18]. The high concentration of BAP would otherwise inhibit organogenesis in the genus *Piper* [12, 17].

Multiplication of red betel in vitro plantlets on MS supplemented with TDZ obtained the highest mean number of shoots (2.6 ± 0.3) and leaves (13.8 ± 4.9) on media supplemented with 0.01 mgL⁻¹ TDZ. The highest mean of shoot lengths (1.7 ± 0.7 cm) was obtained on media supplemented 0.3 mgL⁻¹ TDZ. Tukey test at 5% level proved that the mean number of shoots on MS (control) was not significantly different from the mean number of shoots on MS supplemented with 0.01 mgL⁻¹ TDZ, 0.03 mgL⁻¹ TDZ, 0.1 mgL⁻¹ TDZ, and 0.3 mgL⁻¹ TDZ (Table 3; Figure 1d)

Table 3 Mean Number of Shoots, Shoot lengths, and Leaves on TDZ Media at 12th week

Concentration of TDZ (mg ⁻¹ L)	Mean Number of Shoots (mean ± SD) ^a	Mean of Shoot lengths (cm) (mean ± SD) ^a	Mean Number of Leaves (mean ± SD) ^a
0.01	2.6 ^a ± 0.8	1.5 ^a ± 0.17	13.8 ^b ± 4.9
0.03	1.6 ^a ± 0.9	1.6 ^a ± 1.0	6.10 ^a ± 2.3
0.10	1.8 ^a ± 1.1	1.6 ^a ± 0.8	5.90 ^a ± 3.4
0.30	2.2 ^a ± 2.3	1.7 ^b ± 0.7	5.10 ^a ± 0.3

^a Means followed by same letters are not significantly different at p ≤ 0.05 according to Tukey multiple range test

TDZ is a kind of diphenyl urea plant growth regulator which can directly triggers plant growth and shoots multiplication in a similar physiological way as the N- substituted natural cytokinins. TDZ can also induce the synthesis and accumulation of endogenous cytokinins that trigger bud formation [21]. The highest mean number of shoots of red betel was obtained on media supplemented with low concentration of TDZ, as in *Scrophularia takesimensis*

[22], *Stevia rebaudiana* [21] and *Nyctanthes arbor* [23].

39 shoots multiplication of red betel plantlets 10 achieved on media supplemented with TDZ. The highest mean number of leaves was obtained on media not supplemented with TDZ. It is due to the high concentration of endogenous cytokinins in red betel plant. The high concentration of cytokinins in the media, such as TDZ, therefore will inhibit the formation of leaves [11] because it accumulates endogenous cytokinins [21]. Plant growth regulators can affect plant's growth even if applied in low concentrations. TDZ concentration and duration of application are important factors in organogenesis [24]. 23

Multiplication of red betel in vitro plantlets on MS supplemented with TDZ and NAA obtained 3 highest mean number of shoots (3.4 ± 1.19) on media supplemented with 0.03 mgL^{-1} TDZ and 0.5 mgL^{-1} NAA. The lowest mean number of shoots (2.0 ± 0.7) was obtained on MS supplemented with 0.5 mgL^{-1} NAA (Table 4; Figure 1e)

Table 4 Mean Number of Shoots, Shoot Lengths, and Leaves on TDZ Combined with NAA Media at 12th week

Concentration of (mg ⁻¹ L)	Mean Number of (mean \pm SD) *	Mean of Shoot lengths (mean \pm SD) *	Mean of Number of (mean \pm SD) *
0.01 + 0.5	$2.2^a \pm 0.8$	$1.4^a \pm 0.6$	$5.6^a \pm 1.5$
0.03 + 0.5	$3.4^a \pm 1.7$	$1.8^a \pm 0.3$	$5.7^a \pm 1.6$
0.05 + 0.5	$2.8^a \pm 0.4$	$2.3^a \pm 1.4$	$4.6^a \pm 1.9$
0.00 + 1	$2.0^a \pm 0.7$	$1.5^a \pm 0.9$	$6.1^a \pm 3.4$

* Means followed by same letters are not significantly different at $p \leq 0.05$ according to Tukey multiple range test

Multiplication of red betel plantlets was performed on media supplemented with TDZ and NAA. Media supplemented with cytokinins and auxin can produce shoots more effectively than media supplemented with TDZ solely [25]. The balancing concentration of cytokinins and auxins is a crucial factor for the growth and development of plants. Therefore, both type of plant growth regulators are needed for plant morphogenesis. *H. car 29*ensis [26], *A. cariensis* [27] also had the highest number of shoots on media supplemented with TDZ and NAA rather than on media supplemented with TDZ solely.

3.3 Root Induction

Red betel shoots were successfully rooted after four weeks. The percentage of red betel plantlets successfully rooted on MS (control), MS supplemented with 0.5 mgL^{-1} NAA, and MS supplemented with 1.5 mgL^{-1} NAA was 100%. Meanwhile, the percentage of red betel plantlets rooted on MS supplemented with 1 mgL^{-1} NAA was 90%. The highest mean number of roots was obtained on MS supplemented 0.5 mgL^{-1} NAA, with 22.8 ± 9.7 roots (Table 5).

Table 5 Mean Number of Roots and Percentage of Rooted Plantlets on Root Induction Media at 12th week

Concentration of NAA (mg ⁻¹ L)	Mean Number of Roots (mean \pm SD) *	Rooting (%)
0	$14.6^a \pm 7.4$	100
0.5	$22.8^a \pm 9.7$	100
1	$14.1^a \pm 8.1$	90
1.5	$22.1^a \pm 9.8$	100

* Means followed by same letters are not significantly different at $p \leq 0.05$ according to Tukey multiple range test

Root induction of *Cinchona succirubra* [28] and *Celastrus paniculatus* [29] had been achieved on media supplemented with NAA. Interestingly, roots of red betel plantlets were able to be induced on media without supplementation NAA. This was probably because red betel already contained endogenous auxin Indole Acetic Acid (IAA) that could be translocated to the roots [18]. IAA is biosynthesized in the apical meristem and then translocated basipetally to induce roots formation. NAA is an auxin-type growth regulator that regulates the formation of roots [18]. In this research, NAA had no significant effect on the formation of roots of red betel plantlets.

3.4 Acclimatization of Plantlets

The in vitro propagated plantlets were acclimatized for four weeks in media containing husk : compost 1 : 1 (Figure 1f), and then transferred to the greenhouse for post-acclimatization in media containing husk : soil : compost 1 : 1 : 1 (Figure 1g). Table 6 shows the viable percentage of red betel plants after four weeks being acclimatized and four weeks being post-acclimatized. The highest survival rates of red betel plants during acclimatization and post-acclimatization were 88% and 100%, respectively (Table 6).

Table 6 Survival Rates of Red Betel Plants during Acclimatization and Post-Acclimatization at 4th week

Root Induction Media (mg ⁻¹ L NAA)	Acclimatization (%)	Post-Acclimatization (%)
0	88 (8/9)	100 (8/8)
0.5	88 (8/9)	100 (8/8)
1	78 (7/9)	100 (7/7)
1.5	88 (8/9)	100 (8/8)

During acclimatization, red betel plants had a high mortality rate, due to the instable environmental conditions [10]. Besides, the low viability in acclimatization was probably caused by the death of root cells which therefore interfered the acquisition of nutrition and water from planting media. Plants in the

genus *Piper* had various survival rates during acclimatization, such as *Piper longum* (70%) [17], and *Piper auritum* Khun (100%) [30]. Survival rate of a plant in the acclimatization period was also influenced by the number of roots. Auxin which was given in the in vitro root induction media affected the physiological response of plant growth and development because auxin interacted with cytokinins which were already synthesized in root meristem. Therefore, there was a balance concentration of auxins and cytokinins which was beneficial for plantlet growth. Red betel plants had a survival rate of 100% during post-acclimatization because they had been adapted to environmental conditions during acclimatization




Figure 1 Micropropagation and Acclimatization of Red Betel (*Piper crocatum* Ruiz and Pav.)

a Shoots Induction of in vitro red betel on MS supplemented with 3 mgL⁻¹ BAP, **b** Shoots Initiation of in vitro red betel on MS supplemented with 3 mgL⁻¹ BAP, **c** Shoots multiplication of in vitro red betel on MS supplemented with 1 mgL⁻¹ BAP, **d** Multiplication of red betel plantlet on MS supplemented with 1 mgL⁻¹ TDZ, **e** Shoots multiplication of in vitro red betel on MS supplemented with 0.03 mgL⁻¹ TDZ and 0.05 mgL⁻¹ NAA, **f** Acclimatization of red betel at first month, **g** post-acclimatization of red betel at third months

4.0 CONCLUSION

Single nodes of red betel explant had been induced and shoots had been initiated on MS supplemented with 3 mgL⁻¹ BAP. Shoot multiplication was optimized on MS supplemented with 1 mgL⁻¹ BAP. All of the plantlets had been induced to produce roots. The best mean number of roots had been successfully achieved on MS media supplemented with 0.5 mgL⁻¹ NAA. The highest percentage of rooted plantlets was 100%. The highest survival rates of red betel during acclimatization and post acclimatization were 88% and 100%, respectively.

15

Acknowledgement

The authors would like to thank The Directorate

General of Higher Education, Ministry of National Education, Indonesia for competitive grant project funding and Prof. Ika Mariska for reviewing this paper.

References

- [1] Tupas, L., Popp, B. And Karl, D., 1994. Dissolved Organic Carbon In Oligotrophic Waters: Experiments On Sample Preservation, Storage And Analysis. *Marine Chemistry*. 45: 207-216.
- [2] Wicaksono, B. D., Handoko, Y. A., Arung, E. T., Kusuma, I.W., Yulia, D., Pancaputra, A. N. And Sandra, F. 2009. Antiproliferative Effect of the Methanol Extract of Piper Crocatum Ruiz and Pav. Leaves on Human Breast (T47D) Cells In-Vitro. *Tropical Journal of Pharmaceutical Research*. 4: 345-352.
- [3] Safithri, M. And Fahma. 2005. Potensi Rebusan Daun Sirih Merah (*Piper Crocatum* Ruiz and Pav.) Sebagai Senyawa Antihiperlipidemia Pada Tikus Putih Galur Sprague Dawley. *Hayati Journal of Bioscience*. 15: 45-48
- [4] Fitriyani, A., Winarti, L., Muslicha, S. And Nuri. 2011. Uji Antiinflamasi Ekstrak Methanol Daun Sirih Merah (*Piper Crocatum* Ruiz And Pav.) Pada Tikus Putih. *Majalah Obat Tradisional*. 16: 34-42.
- [5] Juliantina, F. R., Citra, D. A. C., Nirwani, B., Numasitoh, T. And Bowo, E. T. 2009. Manfaat Sirih Merah (*Piper Crocatum* Ruizand Pav.) Sebagai Agen Anti Bakteri Gram Positif Dan Negatif. *Jurnal Kedokteran Dan Kesehatan Indonesia*. 1-10.
- [6] Zubier, F., Bramono, K., Widaty, S., Nilasari, H., Louisa, M. And Rosana, Y. 2010. Efikasi Sabun Ekstrak Sirih Merah Dalam Mengurangi Gejala Keputihan Fisiologi. *Majalah Kedokteran Indonesia*. 40: 9-14.
- [7] Alfarabi, M., Bintang, M., Suryani and Safitri, M. 2010. The Comparatibe Ability of Antioxidant Activity Of Piper Crocatum Ruiz And Pav. In Inhibiting Fatty Acid Oxidation And Free Radical. *Hayati Journal of Bioscience*. 17: 201-204.
- [8] Reveny, J. 2011. Daya Antimikroba Ekstrak Dan Fraksi Daun Sirih Merah (*Piper Betle* Linn.). *Jurnal Ilmu Dasar*. 12: 6-12.
- [9] Werdhany, W. I., Marton, A. And Setyorini, W. 2008. Sirih Merah. Yogyakarta: Balai Pengkajian Teknologi Pertanian Yogyakarta.
- [10] Prastowo, N. H., Roshetko, J. M., Maurung, G. E. S., Nugraha E., Tukan, J. M. And Harum, F. 2006. Teknik Pembibitan Dan Perbanyakan Vegetatif Tanaman Buah. Bogor: ICRAF.
- [10] Gunawan, L. W. 1987. Teknik Kultur Jaringan. Bogor: 1987.
- [11] George, E. F. And Sherrington, L. W. 1984. Plant Propagation By Tissue Culture Handbook And Directory Of Comercial Laboratories. England: Exegetics Ltd.
- [12] Hussain, A., Nazand, S. And Nazir, H. 2011. Tissue Culture Of Black Pepper (*Piper Nigrum* L.) In Pakistan. *Pakistan Journal Of Botany*. 2: 1069-1078.
- [13] Swarna, J. And Ravindhran, R. 2012. In Vitro Propagation And Assessment Of Genetic Integrity Of *Talinum Triangulare* (Jacq) Wild: A Valuable Medicinal Herb. *Acta Physiologiae Plantarum*. 34: 1987-1996.
- [14] Nirwan And Aziz, A. 2006. Multiplikasi Dan Pigmentasi Antosianin Daun Dewa (*Gynura Pseudochina* (L) DC) In Vitro. *Bulletin Agronomi*. 34: 112-118.
- [15] Ade, R. And Rai, M. 2011. Multiple Shoot Information In *Gloriosa Superba*: A Rare And Endangered Indian Medicinal Plant. *Bioscience*. 3: 68-72.
- [16] Vijayakumar, M., Vijayakumar, R. And Stephen, R. 2010. In Vitro Propagation Of *Bacopa Monnieri*-A Multipurpose Medical Plant. *Indian Journal Science And Technology*. 3: 782-789.
- [17] Parida, R. And Dhal, Y. 2011. A Study On The Micro-Propagation And Antioxidant Activity Of *Piper Longum*

- (An Important Medical Plant). *Journal Of Medicinal Plant Research*. 5: 6991-6994.
- [18] Wattimena, G. A. Bioteknologi Tanaman Bogor: Departemen Pendidikan Dan Kebudayaan Direktorat Jenderal Pendidikan Tinggi Pusat Antar Universitas Bioteknologi Institut Pertanian Bogor.
- [19] Saha, S., Kader, A., Sengupta, C. And Ghosh, P. 2011. In Vitro Propagation Of *Ocimum Gratissimum* L. (Lamiaceae) And Its Evaluation Of Genetic Fidelity RAPD Marker. *American Journal Of Plant Science*. 3: 64-74.
- [20] Sharmar, J., Khan, S. And Varman, R. 2012. In Vitro Culture Establishment And Shoot Induction Of *Mallotus Philippensis* (Lam.) M. ARG. *Pharmacie Globale International Journal Of Comprehensive Pharmacy*. 3: 1-5.
- [21] Lata, H., Chandra, S., Wang, Y. H., Raman, V. And Khan, I. A. 2013. TDZ-Induced High Frequency Plant Regeneration Through Direct Shoot Organogenesis In *Stevia Rebaudiana* Bertoni: An Important Medicinal Plant And A Natural Sweetener. *American Journal Of Sciences*. 4: 117-128.
- [22] Sivanesan, I. S., Limand, M. Y. And Jeong, B. R. 2012. Micropropagation And Greenhouse Cultivation Of *Scrophularia Takesimensis* Nakai, A Rare Endemic Medicinal Plant. *Pakistan Journal Of Botany*. 44: 1657-1662.
- [23] Amsar Siddiqueet, I., Anis, M. And Jahan, A. 2006. Rapid Multiplication Of *Nyctanthes Arbor-Trifis* L. Through In Vitro Axillary Shoots Proliferation. *World Journal Of Agricultural Science*. 2: 1881-192.
- [24] Malik, S., Sharma, S., Sharmaand, M. And Ahuja, P.S. 2010. Direct Shoot Regeneration From Intact Leaves Of *Arnebia Euchroma* (Royle). *Cell Biology International*. 34: 537-542.
- [25] Schulze, J. 2007. *Improvements In Cereal Tissue Culture By Thiadiazuron: A Review*. Global Science Book. 64-74.
- [26] He, S. S., Liu, C. Z. And Saxena, P.K. 2007. *Plant Regeneration Of An Endangered Medicinal Plant Hydrastis Canadensis* L. Elsevier. 113: 82-86.
- [27] Erisen, S., Atalay, E., Yorgancilar, M. 2011. The Effect Of Thiadiazuron On The In Vitro Shoot Development Of Endemic *Antragalus Cariensis* In Turkey. *Turkey J Bot*. 35: 521-526.
- [28] Riyadi, I. And Tahardi, J.S. 2005. Pengaruh NAA Dan IBA Terhadap Pertumbuhan Dan Perkembangan Tunas Kina [*Cinchona Succirubra*]. *Jurnal Bioteknologi Pertanian*. 10: 45-50.
- [29] Phulwaria, M., Rai, M. K., Patel, A. K. Kataria, V. And Shekhawat, N. S. 2012. A Genetically Stable Rooting Protocol For Propagating A Threatened Medicinal Plant- *Cleistanthus paniculatus*. *Aob Plant*. 1-9.
- [30] Dominguez, F., Lozoya, X. And Simon, J. 2006. Tissue Culture Regeneration Of A Medical Plant From Mexico: *Piper Auritum* Kunth. *Hortscience*. 41: 207-209.

(Check) 9 - 2016 Jurnal Teknologi(AR2BIO) Nesti Sianipar - Induction, Multiplication and Acclimatization of Red Betel

ORIGINALITY REPORT

%**22**

SIMILARITY INDEX

%**12**

INTERNET SOURCES

%**18**

PUBLICATIONS

%**10**

STUDENT PAPERS

PRIMARY SOURCES

1

ifrj.upm.edu.my

Internet Source

%**2**

2

Submitted to Universiti Putra Malaysia

Student Paper

%**2**

3

Submitted to Higher Education Commission
Pakistan

Student Paper

%**1**

4

www.jurnalteknologi.utm.my

Internet Source

%**1**

5

Ahmed, Sheikh Rashel, Md Moniruzzaman
Shohag Howlader, Pijush Sutradhar, and
Sabina Yasmin. "An efficient protocol for in
vitro regeneration of Stevia rebaudiana", Asian
Journal of Medical and Biological Research,
2016.

Publication

%**1**

6

www.banglajol.info

Internet Source

%**1**

7	Hassan, AKM Sayeed, Farhana Afroz, Laila Shamroze Bari, John Liton Munshi, Miskat Ara Akhter Jahan, and Rahima Khatun. "Micropropagation of <i>Eclipta alba</i> (Linn.) Hassk-a Valuable Medicinal Herb", Bangladesh Journal of Scientific and Industrial Research, 2008. Publication	%1
8	www.rroj.com Internet Source	%1
9	www.cibtech.org Internet Source	%1
10	YÜCESAN, Bahtiyar Buhara, MOHAMMED, Aliyu, ARSLAN, Merve and GÜREL, Ekrem. "Clonal propagation and synthetic seed production from nodal segments of Cape gooseberry (<i>Physalis peruviana</i> L.), a tropical fruit plant", TÜBİTAK, 2015. Publication	%1
11	www.unsjournals.com Internet Source	%1
12	Junli Wang. "An efficient callus proliferation protocol and rhaponticin accumulation of <i>Rheum franzenbachii</i> Munt., a medicinal plant", Journal of Plant Biochemistry and Biotechnology, 04/27/2011 Publication	%1

13	Huq, Amdadul, Shahina Akter, Shahina Islam, and Salim Khan. "In vitro micropropagation of pointed gourd (<i>Trichosanthes dioica</i> Roxb.) from shoot tip and nodal segment", Bangladesh Journal of Scientific and Industrial Research, 2012.	%1
----	---	----

Publication

14	Md. Mahabubur Rahman. "Mass propagation of <i>Solanum surattense</i> Bum. using direct adventitious shoot organogenesis from internode", Acta agriculturae Slovenica, 03/01/2011	%1
----	--	----

Publication

15	ajouronline.com	%1
----	--	----

Internet Source

16	Biotechnological strategies for the conservation of medicinal and ornamental climbers, 2016.	%1
----	--	----

Publication

17	search.oecd.org	<%1
----	--	-----

Internet Source

18	CHALAGERI, Gururaj and BABU, Uddagiri Venkanna. "In vitro plant regeneration via petiole callus of <i>Viola patrinii</i> and genetic fidelity assessment using RAPD markers", TÜBİTAK, 2012.	<%1
----	--	-----

Publication

20

Ahmed, Madi Waheed Al-Mayahi. "Thidiazuron-induced in vitro bud organogenesis of the date palm (*Phoenix dactylifera* L.) CV. Hillawi", AFRICAN JOURNAL OF BIOTECHNOLOGY, 2014.

Publication

<% 1

21

Su Tiing Sai. "In vitro propagation of *Typhonium flagelliforme* (Lodd) blume", In Vitro Cellular & Developmental Biology - Plant, 09/2000

Publication

<% 1

22

Dessalegn, Obssi, Kassahun Bantte, and Mulugeta Diro. "Effect of Different Combinations of Plant Growth Regulators on in vitro Propagation of Yam (*Dioscorea* Species)", Journal of Applied Biotechnology, 2015.

Publication

<% 1

23

Khatun, MK, MS Haque, S Islam, and KM Nasiruddin. "In vitro Regeneration of Mungbean (*Vigna radiata* L.) from Different Explants", Progressive Agriculture, 2013.

Publication

<% 1

24

Karim, M Rezaul, MA Malek, Sajia Rahman, M Al-Amin, and M Ruhul Amin. "In vitro propagation of banana", Bangladesh Journal of

<% 1

-
- 25 Mahipal S. Shekhawat, M. Manokari, J. Revathi. "<div>In vitro propagation and ex vitro rooting of Aerva lanata (L.) Juss. ex Schult.: a rare medicinal plant", Indian Journal of Plant Physiology, 2016

Publication

-
- 26 Meena K. Cheruvathur. "An efficient plant regeneration system through callus for Pseudarthria viscida (L.) Wright and Arn., a rare ethnomedicinal herb", Physiology and Molecular Biology of Plants, 10/04/2011

Publication

-
- 27 Submitted to Jawaharlal Nehru Technological University

Student Paper

-
- 28 RIZWAN, M.; SHARMA, R.; SONI, P.; GUPTA, N. K. and SINGH, G.. "REGENERATION PROTOCOL FOR CHILLI (CAPSICUM ANNUUM L.) VARIETY MATHANIA", Journal of Cell & Tissue Research, 2013.

Publication

-
- 29 academicjournals.org

Internet Source

-
- 30 "Research Reports on Botany from Gyeongsang National University Provide New

31

Biotechnology for Medicinal Plants, 2013.

Publication

<% 1

32

eprints.undip.ac.id

Internet Source

<% 1

33

Junliang Zhao. "Transformation of modified cowpea trypsin inhibitor gene and anti-bacterial peptide gene in Brassica pekinensis protoplasts mediated by Agrobacterium tumefaciens", Euphytica, 06/2006

Publication

<% 1

34

Sandra Gonçalves. "An efficient in vitro propagation protocol for Pinguicula lusitanica, a rare insectivorous plant", Plant Cell Tissue and Organ Culture, 11/2008

Publication

<% 1

35

M. Evtimova. "Flax Improvement by Biotechnology Means", Journal of Natural Fibers, 08/31/2005

Publication

<% 1

36

Hartini, YS, S Wahyuono, S Widyarini, and A Yuswanto. "In vivo immunomodulatory effect and histopathological features of mouse liver and kidney treated with neolignans isolated

<% 1

from red betel (*Piper crocatum* Ruiz & Pav)
leaf", Tropical Journal of Pharmaceutical
Research, 2014.

Publication

- 37 Mng'omba, S.A.. "Repeated exposure of jacket
plum (*Pappea capensis*) micro-cuttings to
indole-3-butyric acid (IBA) improved in vitro
rooting capacity", South African Journal of
Botany, 200704

Publication

<% 1

- 38 V. A. Parthasarathy. "Black Pepper: Botany and
Horticulture", Horticultural Reviews,
01/01/2007

Publication

<% 1

- 39 Shinde, Amit N.. "Induced High Frequency
Shoot Regeneration and Enhanced Isoflavones
Production in *Psoralea corylifolia*", Records of
Natural Products/13076167, 20090101

Publication

<% 1

- 40 Biotechnology in Agriculture and Forestry,
1992.

Publication

<% 1

EXCLUDE QUOTES ON
EXCLUDE ON
BIBLIOGRAPHY

EXCLUDE MATCHES < 5 WORDS